

AD-A036 724

INDIANA UNIV BLOOMINGTON DEPT OF MATHEMATICS F/G 12/1
LOCAL MAXIMA OF THE SAMPLE FUNCTIONS OF THE N-PARAMETER BESSEL --ETC(U)
1977 M L PURI, L T TRAN AF-AFOSR-2927-76

UNCLASSIFIED

AFOSR-TR-77-0111

NL

1 of 1
ADA036724




END

DATE
FILMED

4-77

1 OF 1

ADA036724



24

97

AD A 036724

* Work supported by the Air Force Office of Scientific Research, A.F.S.C.U.S.A.F., Grant No. AFOSR76-2000. 2927
Reproduction in whole or part permitted for any purpose of the United States Government.

[illegible]

N -parameter Wiener process, the components being independent. Write $W = W^{(N,d)}$ for simplicity, and denote the i th component of W by W^i . Define the N -parameter Bessel process associated with W by

$$(1.1) \quad B_t = \left[\sum_{i=1}^d (W_t^i)^2 \right]^{\frac{1}{2}}$$

It is shown that almost every sample function of B_t has a local maxima. Furthermore some properties related to the local maxima of B_t are investigated.

As in Orey and Pruitt** (1973), our parameter space is R_+^N , that is the set of $t \in R^N$ with all components non-negative. When dealing with a point t in the parameter space we sometimes write $t = \langle t_1, \dots, t_N \rangle$ or simply $\langle t_i \rangle$. In case all $t_i = 0$, we write $t = \langle 0 \rangle$. For $s = \langle s_i \rangle$ and $t = \langle t_i \rangle$ with $s_i \leq t_i$, the interval $\bigcap_{i=1}^N [s_i, t_i]$ is denoted by $\Delta(s, t)$, and by $\Delta(t)$ in case $s = \langle 0 \rangle$. Denote by $S(s, t)$, the symmetric difference of $\Delta(s)$ and $\Delta(t)$. Then it is easy to check that if $s, t \in R_+^N$, the variance of $W^i(t) - W^i(s)$ is $|S(s, t)|$ where $|\cdot|$ denotes the N -dimensional Lebesgue measure. Furthermore,

** Wherever possible, we shall use the notation of Orey and Pruitt (1973).

W has continuous sample functions and independent increments. We denote the increment of W over $\Delta(s,t)$ by $W(\Delta(s,t))$. For further information on W , the reader is referred to Kitagawa (1951), Chentsov (1956), Yeh (1960, 1963a, 1963b), Park, C. (1969), Park, W. J. (1970), Zimmerman (1972) and Orey and Pruitt (1973).

Definition 1.1. The sample function $B(\cdot, \omega)$ has a local maxima at s if there exists an open set O containing s such that $O \subset R_+^N$ and $B(t, \omega) \leq B(s, \omega)$ for all $t \in O$.

We shall need the Orey-Pruitt analogue of the familiar zero-one law. Let C_n be the class of time intervals in R_+^N with vertices of the form $\langle k_i 2^{-n} \rangle$, k_i nonnegative integers, and having all sides of equal length, and for $n > 0$ each member of C_n is to be a subcube of one in C_0 . Let $C_\infty = \bigcup_{n=0}^{\infty} C_n$, and $\mathcal{F}_n = \mathcal{B}(W(\Delta), \Delta \in C_n)$, $\mathcal{F}_\infty = \bigvee_{n=0}^{\infty} \mathcal{F}_n$. Thus \mathcal{F}_n is the Borel field generated by the indicated class of random variables and \mathcal{F}_∞ is the smallest Borel field including all \mathcal{F}_n . For a subset D of R_+^N , we put $C_n(D) = \{\Delta \in C_n; \Delta \subseteq D\}$, $\mathcal{F}_n(D) = \mathcal{B}\{W(\Delta); \Delta \in C_n(D)\}$, $\mathcal{F}_\infty(D) = \bigvee_{n=0}^{\infty} \mathcal{F}_n(D)$ then we have the following lemma.

Lemma 1.1 (Orey-Pruitt (1973)). Let $D_m \subset R_+^N$, $m = 1, 2, \dots$,

with $D_m \neq \emptyset$. If $A \in \mathcal{F}_\infty(D_m)$ for every m , then $P(A) = \{0,1\}$.

Lemma 1.2. Let φ be a nonnegative, nondecreasing,
continuous function defined for large arguments. Then for
almost all w there is an $\varepsilon(w)$ such that for all intervals
 $\Delta(s,t)$ with $\Delta(s,t) \subset \Delta(\langle 1 \rangle)$ and $|\Delta(s,t)| < \varepsilon(w)$,

$$|W(\Delta(s,t))| < |\Delta(s,t)|^{\frac{1}{2}\varphi(|\Delta(s,t)|^{-1})}$$

if and only if

$$\int_0^\infty \log \varepsilon^{3N+d/2-2} e^{-\varphi^2(\varepsilon)/2} d\varepsilon$$

converges.

For the proof of this lemma, see Orey and Pruitt
 (1973, page 147).

2. Local Maxima. In this section we prove the main theorem
 dealing with the existence of the local maxima of the sample
 functions of the Bessel process B_t .

Theorem 2.1. For almost all sample functions of the Bessel process B_t defined in (1.1), there exists a local maxima.

Proof. Let s be the center of the unit interval U , and let $C_n \subset U$ be a cube with center at s , sides parallel to the coordinate axes and equal to a_n . Let u^n and v^n be the smallest and the largest vertex of C_n i.e. closest and farthest from the origin $\langle 0 \rangle$. Pick C_n with $\min(u_1^n, \dots, u_N^n) > \frac{1}{4}$.

Consider two points s^{nk} and v^{nk} of R_+^N determined by $s_k^{nk} = \frac{1}{2}$, $v_k^{nk} = v_k^n$, $s_j^{nk} = v_j^{nk} = u_j^n$ for $j \neq k$ where $1 \leq j < N$.

Let $\Delta(r, \ell)$ be any interval in U with at least two sides smaller than a_n . Define

$$A_{ni} = \bigcap_{k=1}^N [W^i(s^{nk}) - W^i(u) > 2a_n^{\frac{1}{2}}, W^i(s^{nk}) - W^i(v^{nk}) > 2a_n^{\frac{1}{2}}]$$

$$B_{ni} = \bigcap_{k=1}^N [W^i(s^{nk}) - W^i(u) < -2a_n^{\frac{1}{2}}, W^i(s^{nk}) - W^i(v^{nk}) < -2a_n^{\frac{1}{2}}]$$

$$C_{ni} = [\inf_{t \in C_n} W_t^i \geq 0],$$

$$E_{ni} = [\sup_{t \in C_n} W_t^i \leq 0],$$

$$F_{ni} = [\sup_{r, \ell \in U} |W^i(\Delta(r, \ell))| < (2^{N-1} - 1)^{-1} a_n^{\frac{1}{2}}]$$

The variables $W^i(s^{nk}) - W^i(u)$, $W^i(s^{nk}) - W^i(v^{nk})$ are normally distributed with mean 0 and variances greater than $4^{-N+\frac{1}{2}}a_n$. Thus $P(A_{ni}) > \beta$ for some constant β .

Let $\{a_n\}$ be a sequence of positive numbers with $a_n \downarrow 0$ and let D_n be the interior of $S(u^n, v^n)$. Clearly $D_n \downarrow \emptyset$ as $a_n \downarrow 0$. Observe that the event $[A_{ni} \text{ infinitely often}] \in \mathcal{F}_\infty(D_n)$. Thus, from Lemma 1.1, it follows that

$$P(A_{ni} \text{ infinitely often}) = 1$$

Analogously,

$$P(B_{ni} \text{ infinitely often}) = 1.$$

Let $\eta > 0$. Then by Lemma 1.2, or by the continuity of the sample functions of W ,

$$P\left[\bigcap_{n=n_0}^{\infty} C_{ni} \mid W_s^i > \eta\right] \rightarrow 1 \text{ as } n_0 \rightarrow \infty$$

Furthermore,

$$\begin{aligned} P[A_{ni} C_{ni} \text{ infinitely often}] &\geq P[(A_{ni} \text{ infinitely often}) \bigcap_{n=n_0}^{\infty} C_{ni}] \\ &\geq P\left[\bigcap_{n=n_0}^{\infty} C_{ni}\right] \\ &\geq P\left[\bigcap_{n=n_0}^{\infty} C_{ni} [W_s^i > \eta]\right] \end{aligned}$$

Since $P[W_s^i > \eta]$ converges to $\frac{1}{2}$ as $\eta \rightarrow 0$, by picking η small enough and then n_0 large enough, the probability of the last event can be made as close to $\frac{1}{2}$ as desired.

It is now clear that

$$P[(A_{ni}C_{ni} \cup B_{ni}E_{ni}) \text{ i.o.}] = 1.$$

Also, by Lemma 1.2 and by the independence of the components of W

$$P[\bigcap_{i=1}^d (A_{ni}C_{ni}F_{ni} \cup B_{ni}E_{ni}F_{ni}) \text{ i.o.}] = 1$$

Let $t \in \partial C_n$ where ∂C_n is the boundary of C_n , and let

$$p^t = \langle \sigma(t_1), \dots, \sigma(t_N) \rangle$$

where

$$\sigma(t_i) = \begin{cases} t_i & \text{if } u_i^n < t_i < v_i^n \\ s_i & \text{otherwise} \end{cases}$$

Observe that p^t lies in the interior of C_n . Now consider events of the form

$$\bigcap_{i=1}^d \wedge_{ni} \text{ where } \wedge_{ni} = A_{ni}C_{ni}F_{ni} \text{ or } B_{ni}E_{ni}F_{ni}$$

and note that there are 2^d events of this form.

We now claim that

$$(2.1) \quad P[\sup_{t \in C_n^0} B_t > \sup_{t \in \partial C_n} B_t \mid \bigcap_{i=1}^d A_{ni}C_{ni}F_{ni}] = 1$$

A slight variation of the proof along the same lines can be applied to $2^d - 1$ other events. To prove (2.1), let

$$(2.2) \quad G_n = \bigcap_{i=1}^d A_{ni} C_{ni} F_{ni}$$

$$H_n = \bigcap_{t \in \partial C_n} \left\{ \bigcap_{i=1}^d [W^i(p^t) - W^i(t) > a_n^{\frac{1}{2}}] \right\}.$$

Now

$$W^i(p^t) - W^i(t) \cong W^i(\langle \sigma(t_1), t_2, \dots, t_N \rangle)$$

$$-W^i(\langle t_1, t_2, \dots, t_N \rangle) + W^i(\langle \sigma(t_1), \sigma(t_2), t_3, \dots, t_N \rangle)$$

$$-W^i(\langle \sigma(t_1), t_2, \dots, t_N \rangle) + \dots + W^i(\langle \sigma(t_1), \sigma(t_2), \dots, \sigma(t_n) \rangle)$$

$$-W^i(\langle \sigma(t_1), \dots, \sigma(t_{N-1}), t_N \rangle).$$

Consider the random variable

$$W^i(\langle \sigma(t_1), \dots, \sigma(t_{j-1}), \sigma(t_j), t_{j+1}, \dots, t_N \rangle)$$

(2.3)

$$-W^i(\langle \sigma(t_1), \dots, \sigma(t_{j-1}), t_j, t_{j+1}, \dots, t_N \rangle).$$

The variance of this variable is equal to

$$(u_1^n + e_1^n) \dots (u_{j-1}^n + e_{j-1}^n) |\sigma(t_j) - t_j| (u_{j+1}^n + e_{j+1}^n) \dots (u_N^n + e_N^n)$$

where $0 \leq e_i^n \leq a_n$. It is now easy to see that (2.3) is equal to

$$W^i(\langle u_1^n, \dots, u_{j-1}^n, \sigma(t_j), u_{j+1}^n, \dots, u_N^n \rangle)$$

$$-W^i(\langle u_1^n, \dots, u_{j-1}^n, t_j, u_{j+1}^n, \dots, u_N^n \rangle) + L^i,$$

where L^i can be decomposed into no more than $2^{N-1} - 1$ normally distributed random variables, such that the mean of each of these random variables is zero and the variance of each is equal to the N -dimensional Lebesgue measure of an interval in U with at least two sides smaller than a_n . Since $t \in \partial C_n$, t_j is equal to u_j^n or v_j^n and $\sigma(t_j) = \frac{1}{2}$ for some $1 \leq j \leq N$. Therefore

$$P(H_n \bigcap_{i=1}^d C_{ni} \mid G_n) = 1$$

However,

$$P[\sup_{t \in C_n} B_t > \sup_{t \in \partial C_n} B_t \mid H_n \bigcap_{i=1}^d C_{ni}] = 1$$

Hence

$$P[\sup_{t \in C_n} B_t > \sup_{t \in \partial C_n} B_t \mid G_n] = 1$$

where G_n is defined in (2.2).

The proof is now completed. Recall that s was picked to be the center of U . Actually, s can be chosen to be any point in U^0 . Therefore, for almost all sample functions of B_t , the set of local maxima is dense in R_+^N .

We shall now investigate some properties of the local maxima of B_t .

Definition 2.1. The sample function $B(\cdot, \omega)$ has a strict local maxima if there exists an open set O containing s such that $0 \in R_+^N$ and $B(t, \omega) < B(s, \omega)$ for all $t \in O$.

We have the following theorem.

Theorem 2.2. For almost every sample function of $\{B_t, t \in R_+^N\}$, all the local maxima are strict and the set of local maxima is countable.

Proof. Let I and J be two disjoint, closed intervals in the interior of R_+^N . We claim that

$$(2.4) \quad P\{\sup_{t \in I} B_t = \sup_{t \in J} B_t\} = 0.$$

Let $I = \Delta(u, v)$, $J = \Delta(s, t)$. Denote the complements of $\Delta(t)$ and $\Delta(v)$ by $[\Delta(t)]'$ and $[\Delta(v)]'$. Since I and J are disjoint intervals, it is clear that

$$[\Delta(u) \cap [\Delta(t)]'] \cup [\Delta(s) \cap [\Delta(v)]']$$

contains a nondegenerate interval, i.e., an interval with positive N -dimensional Lebesgue measure. Let I' be any such interval, and without loss of generality assume that

$$I' \subset \Delta(u) \cap [\Delta(t)]'$$

Consider now

$$\begin{aligned} & P\left[\sup_{t \in I} B_t = \sup_{t \in J} B_t\right] \\ &= P\left[\sup_{t \in I} \left[(W_t^1)^2 + \sum_{i=2}^d (W_t^i)^2\right]^{\frac{1}{2}} = \sup_{t \in J} \left[\sum_{i=1}^d (W_t^i)^2\right]^{\frac{1}{2}}\right]. \end{aligned}$$

Let $W^1(I')$ be the increment of W^1 over I' . Since W^1 has independent increments, for $t \in I$, we can write

$$W_t^1 = W_t^1 - W^1(I') + W^1(I')$$

such that $W^1(I')$ is independent of $W_t^1 - W^1(I')$ for all $t \in I$. Also, $W^1(I')$ is independent of $\sup_{t \in J} \left[\sum_{i=1}^d (W_t^i)^2\right]$ since $I' \subset [\Delta(t)]'$.

Let $X = W^1(I')$, $Y_t = W_t^1 - W^1(I')$. For an arbitrary fixed w , consider

$$\begin{aligned} (2.5) \quad & P\left[\sup_{t \in I} \{(X + Y_t(w))^2 + \sum_{i=2}^d (W_t^i(w))^2\}^{\frac{1}{2}}\right. \\ & \left.= \sup_{t \in J} \left[\sum_{i=1}^d (W_t^i(w))^2\right]^{\frac{1}{2}}\right]. \end{aligned}$$

We shall now show that (2.5) equals zero for a fixed w .

Consider the function $f(x)$, defined by

$$f(x) = \sup_{t \in I} \{(x + Y_t(w))^2 + \sum_{i=2}^d (W_t^i(w))^2\}^{\frac{1}{2}}.$$

Note that $f(x)$ equals the supremum of the distance from the origin of the set D_x in R^d defined by

$$D_x = \{z \in R^d : z_1 = y_t(u) + x, z_2 = w_t^2(u), \dots, z_d = w_t^d(u) \text{ for some } t \in I\}.$$

It is now easy to see that as x varies from $-\infty$ to $+\infty$, the set D_x is translated along a vector parallel to a coordinate axis and so $f(x)$ decreases and then increases as x goes from $-\infty$ to $+\infty$. For a fixed u , (2.5) equals

$$(2.6) \quad P[X = f^{-1}(\sup_{t \in J} \sum_{i=1}^d (w_t^i(u))^2)]$$

where f^{-1} is the inverse of f . It is clear that there are almost 2 values of $f^{-1}(\sup_{t \in J} \sum_{i=1}^d (w_t^i(u))^2)$, and since X is normal random variable, (2.6) equals 0. Thus, for each fixed u , (2.5) equals zero. The proof of (2.4) follows by integrating (2.5) over the probability space.

Consider the set

$$\cap [\sup_{t \in I} B_t \neq \sup_{t \in J} B_t]$$

where the intersection is taken over all intervals I and J with rational least and largest vertices, i.e., the coordinates of u, v, s, t are all rational. This set contains the set of u such that all local maxima of $B(\cdot, u)$ are strict. Clearly, this set has probability one.

Countability of the set of local maxima is a consequence of the following Lemma.

Lemma 2.1. Let f be a continuous, real valued function on \mathbb{R}_+^N with all local maxima strict. Then f has countably many local maxima.

This Lemma is a straightforward generalization of the univariate case, the proof of which can be found in Freedman (1971).

REFERENCES

- [1] Chentsov, N. N. (1956). Wiener random fields depending on several parameters. Dokl. Akad. Nauk. SSSR 106, 607-609.
- [2] Delporte, J. (1966). Fonctions aléatoires de deux variables presque sûrement à échantillons continus sur un domaine rectangulaire borné. Z. Wahrscheinlichkeitstheorie und verw. Gebiete 6, 181-205.
- [3] Freedman, D. (1971). Brownian motion and diffusion. Holden Day, San Francisco.
- [4] Kitagawa, T. (1951). Analysis of variance applied to function spaces. Mem. Fac. Sci. Kyushu Univ. Ser. A. 6, 41-53.
- [5] Orey, S. and Pruitt, W. E. (1973). Sample functions of the N-parameter Wiener process. Ann. Prob. 1, 138-163.
- [6] Park, C. (1969). A generalized Paley-Wiener-Zygmund integral and its applications. Proc. Amer. Math. Soc. 23, 388-400.
- [7] Park, W. J. (1970). A multiparameter Gaussian process. Ann. Math. Statist. 41, 1582-1595.
- [8] Yeh, J. (1960). Wiener measure in a space of functions of two variables. Trans. Amer. Math. Soc. 95, 443-450.
- [9] Yeh, J. (1963a). Cameron-Martin translation theorems in the Wiener space of functions of two variables. Trans. Amer. Math. Soc. 107, 409-420.
- [10] Yeh, J. (1963b). Orthogonal developments of functionals and related theorems in the Wiener space of functions of two variables. Pacific J. Math. 13, 1427-1436.
- [11] Zimmerman, G. (1972). Some sample function properties of the two-parameter Gaussian process. Ann. Math. Statist. 43, 1235-1246.

ADDITIONAL INFO		NOTE SECTION	
NTS		NOTE SECTION	
DOC		BUT SECTION	
UNANNOUNCED			
JUSTIFICATION			
BY		DISTRIBUTION AVAILABILITY CODES	
DIST		AVAIL. AND OR SPECIAL	
A			

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR - 77-0111	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) LOCAL MAXIMA OF THE SAMPLE FUNCTIONS OF THE N-PARAMETER BESSEL PROCESS.		5. TYPE OF REPORT & PERIOD COVERED 9 Interim rept.
7. AUTHOR(s) Madan L. Puri and Lanh Tat Tran		6. PERFORMING ORG. REPORT NUMBER AFOSR 76-2927
9. PERFORMING ORGANIZATION NAME AND ADDRESS Indiana University Department of Mathematics Bloomington, Indiana 47401		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332		12. REPORT DATE 1977
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 15p.		13. NUMBER OF PAGES 14
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		15. SECURITY CLASS. (of this report) UNCLASSIFIED
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) (15) AF-AFOSR-2927-76		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
18. SUPPLEMENTARY NOTES (16) 2304 (17) A5 (18) AFOSR		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) TR-77-0111		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Bessel process local maxima Wiener process independent increments sample functions zero-one law		
21. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is proved that almos every sample function of the N-Parameter Bessel process has a local maxima. In addition, some properties related to the local maxima are investigated.		